

# 2021 Recognition Program

# Achievement Awards

Each of these 7 Achievement Measures must independently be at 85% or higher for the award timeframe in order to receive this award.

1. IV alteplase arrive by 3.5 hour, treat by 4.5 hour
2. Early antithrombotics
3. VTE prophylaxis
4. Antithrombotics
5. Anticoagulation for AFib/AFlutter
6. Smoking cessation
7. Intensive statin therapy

These Performance Achievement Award levels reflect the amount of time for which a hospital must demonstrate compliance to the above eligibility criteria:

**Bronze:** Recognizes performance on 3 consecutive months in a calendar quarter, with a minimum of 30 patients

**Silver:** Recognizes performance on 12 consecutive months, on a January-December time frame


**Gold:** Recognizes performance on 24 consecutive months, on a January-December time frame

# IV alteplase arrive by 3.5 hour, treat by 4.5 hour

Percent of acute ischemic stroke patients who arrive at the hospital within 210 minutes (3.5 hours) of time last known well and for whom IV alteplase was initiated at this hospital within 270 minutes (4.5 hours) of time last known well.

**Documented exclusions or relative exclusions (contraindications or warnings) for not initiating IV thrombolytic in the 0-3 hr treatment window?**  Yes  No

Documented exclusions or relative exclusions (contraindications or warnings) for not initiating IV thrombolytic in the 3-4.5 hr treatment window?  Yes  No



Exclusion Criteria (contraindications) 3-4.5 hr treatment window. Select all that apply:

- C1: Elevated blood pressure (systolic > 185 mm Hg or diastolic > 110 mm Hg) despite treatment
- C2: Recent intracranial or spinal surgery or significant head trauma, or prior stroke in previous 3 months
- C3: History of previous intracranial hemorrhage, intracranial neoplasm, arteriovenous malformation, or aneurysm
- C4: Active internal bleeding
- C5: Acute bleeding diathesis (low platelet count, increased PTT, INR >= 1.7 or use of NOAC)
- C6: Symptoms suggest subarachnoid hemorrhage
- C7: CT demonstrates multilobar infarction (hypodensity >1/3 cerebral hemisphere)
- C8: Arterial puncture at noncompressible site in previous 7 days
- C9: Blood glucose concentration <50 mg/dL (2.7 mmol/L)

Relative Exclusion Criteria (Warnings) 3-4.5 hr treatment window. Select all that apply:

- W1: Care-team unable to determine eligibility
- W2: IV or IA thrombolysis/thrombectomy at an outside hospital prior to arrival
- W3: Life expectancy < 1 year or severe co-morbid illness or CMO on admission
- W4: Pregnancy
- W5: Patient/family refusal
- W7: Stroke severity too mild (non-disabling)
- W8: Recent acute myocardial infarction (within previous 3 months)
- W9: Seizure at onset with postictal residual neurological impairments
- W10: Major surgery or serious trauma within previous 14 days
- W11: Recent gastrointestinal or urinary tract hemorrhage (within previous 21 days)

Additional Relative Exclusion Criteria 3-4.5 hr treatment window. Select all that apply:

- AW1: Age > 80
- AW2: History of both diabetes and prior ischemic stroke
- AW3: Taking an oral anticoagulant regardless of INR
- AW4: Severe stroke (NIHSS > 25)

# Intensive statin therapy

Percentage of Ischemic Stroke and TIA patients who are prescribed high-intensity statin therapy at discharge OR, if > 75 years of age, are prescribed at least moderate- intensity statin therapy at discharge.

<b>Cholesterol Reducing Tx (Select all that apply):</b>	<input type="checkbox"/> None prescribed/ND	<input type="checkbox"/> Statin	<input type="checkbox"/> Niacin	<input type="checkbox"/> Other med
	<input type="checkbox"/> None - contraindicated	<input type="checkbox"/> Fibrate	<input type="checkbox"/> Absorption inhibitor	<input type="checkbox"/> PCSK 9 inhibitor
<b>Statin Medication</b>	<input type="text"/>	<b>Statin Total Daily Dose</b>	<input type="text"/>	
<b>Documented Reason for Not Prescribing Guideline Recommended Dose?</b>	<input type="checkbox"/> Intolerant to moderate (greater than 75yr) or high (less than or equal to 75yr) intensity statin			
	<input type="checkbox"/> No evidence of atherosclerosis (cerebral, coronary, or peripheral vascular disease)			
	<input type="checkbox"/> Other documented reason			
	<input type="checkbox"/> Unknown/ND			

# Quality Awards

At least 5 of these Quality Measures must independently be at 75% or higher for the award time frame in order to receive this add-on award.

1. Dysphagia screen
2. Time to intravenous thrombolytic therapy – 60 min
3. NIHSS reported
4. Stroke education
5. Rehabilitation considered
6. LDL documented

# PHASE III TARGET: STROKE<sup>SM</sup>


HIGHER GOALS FOR GREATER GOOD

## PHASE III RECOGNITION LEVELS

- **Honor Roll:** door-to-needle times within 60 minutes for at least 75 percent of applicable patients
- **Honor Roll Elite:** door-to-needle times within 60 minutes for at least 85 percent of applicable patients
- **Honor Roll Elite Plus:** door-to-needle times within 45 minutes for at least 75 percent of applicable patients and door-to-needle times within 30 minutes for at least 50 percent of applicable patients
- **Honor Roll Advanced Therapy:** door-to-device times in at least 50% of applicable patients within 90 minutes for direct arriving and within 60 minutes for transfers

## ELIGIBILITY

A hospital must currently hold Gold, Silver or Bronze performance achievement status in Get With The Guidelines®-Stroke to be eligible for the Target: Stroke Honor Roll. It must also have DTN/DTD times that meet the criteria above for consecutive applicable patients (minimum six) for at least one calendar quarter for initial awards and four calendar quarters for renewal. .



2 Treatment windows:  
6 or 24 hours



American Heart Association.

Target: Type 2 Diabetes™

## Target: Type 2 Diabetes Honor Roll (can be added-on to Silver, or Gold award)

A composite of all required measures listed below must be at 90% or higher for the award time frame in order to receive this add-on award.

Hospital must have at least 10 patients with a new onset or previous history of diabetes within the patient population.

1. IV alteplase arrive by 3 hour, treat by 4.5 hour for patients with Diabetes
2. Early antithrombotics for patients with Diabetes
3. VTE prophylaxis for patients with Diabetes
4. Antithrombotics for patients with Diabetes
5. Anticoagulant for AFib/Aflutter for patients with Diabetes
6. Smoking cessation for patients with Diabetes
7. Intensive Statin therapy for patients with Diabetes
8. Diabetes treatment
9. Therapeutic Lifestyle Recommendation for Patients with Diabetes