




## Oklahoma State Fact Sheet

Leading Causes of death in Oklahoma in 2010\*



**Oklahoma has the 3rd highest death rate from cardiovascular disease in the country.\*\***

- Heart disease is the No. 1 killer in Oklahoma\*
- Stroke is the No. 5 killer in Oklahoma\*
- 9,426 people in Oklahoma died of heart disease in 2010\*
- 1,980 people in Oklahoma died of stroke in 2010\*

### Heart Disease and Stroke Risk Factors in Oklahoma

	Oklahoma	US
Adults who are current smokers	26.1%	21.1%
Adults who participate in 150+ min of aerobic physical activity per week	44.8%	51.6%
Adults who are overweight or obese+	65.5%	63.5%
Adults who have been told that they have had a heart attack	5.3%	4.4%
Adults who have been told that they have had a stroke	3.4%	2.9%
Adults who have been told that they have angina or coronary heart disease	4.9%	4.1%
Population of adults (18-64) who have some kind of health care coverage	73.4%	78.9%
High school Students who are obese++	16.7%	13%

\* Based on total number of deaths in 2010. Centers for Disease Control and Prevention. WISQARS Leading Cause of Death Reports, 2010.

^ Also known as COPD (Chronic Obstructive Pulmonary Disorder); COPD and allied conditions (including asthma); the term in the ICD/10 is "chronic lower respiratory diseases."

\*\* List includes Puerto Rico and D.C. Based on 2007-09 age-adjusted death rates (ranked from worst to best CVD health). American Heart Association. Heart Disease and Stroke Statistics, 2013 Update: A Report from the American Heart Association. Circulation. Accessed, Jan. 18, 2013.

\*\*\*Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System Survey, 2011.

+ Overweight is defined as having a body mass index (BMI) of 25.0-29.9. Obese is defined as having a body mass index of 30.0 or more.

++ Students who were ≥ 95th percentile for body mass index, by age and sex. Centers for Disease Control and Prevention, Youth Risk Behavior Surveillance System, 2011.