



Post Cardiac Arrest Care – Fact Sheet

Get With The Guidelines®-Resuscitation collects data on resuscitation events from hospitals across the U.S. The data is used to provide participating hospitals with feedback on their resuscitation practice and patient outcomes as well as to develop new evidence-based guidelines for resuscitation.

This fact sheet provides an overview of the five Post-Cardiac Arrest Care (PCAC) measures. These measures were developed by the Get With The Guidelines®-Resuscitation expert leaders to help support hospital processes and aspects of care that are strongly supported by science. With the ultimate goal of improving patient outcomes, the PCAC measures look specifically at targeted temperature management, coronary reperfusion, and optimization of hemodynamics and gas exchange.

MEASURES

- Targeted Temperature Management:
 Percent of adult and pediatric events
 with a cardiac arrest event and return
 of spontaneous circulation (ROSC),
 who are not following commands at
 the time of the initial assessment, in
 whom Targeted Temperature
 Management was utilized.
- Targeted Temperature Distribution: Patients grouped by targeted temperatures
- Door to Cath Lab Times (STEMI): Time from arrival to catheterization lab for adult patients with STEMI (out of hospital events).

- Oxygen Titration: Percent of patients with an arterial blood gas documented with PaO2 maintained at less than 300mmHg within the first 24 hours after ROSC.
- Hypotension Management: Percent of patients with a cardiac arrest event and return of spontaneous circulation (ROSC) with appropriate management of sustained hypotension.